

Seminario de análisis matemático y aplicaciones
Analisi matematikoa eta aplikazioak mintegia

The Pohozaev identity for the fractional Laplacian

XAVIER ROS-OTON

Universitat Politècnica de Catalunya

ABSTRACT: The Pohozaev identity satisfied by solutions of $-\Delta u = f(x, u)$ in Ω , $u = 0$ on $\partial\Omega$, is a classical result which has many applications. For instance, it leads to sharp nonexistence results, to unique continuation properties, monotonicity formulas, or symmetry of solutions.

The aim of this talk is to present the Pohozaev identity for the fractional Laplacian. In a joint work with Joaquim Serra, we establish a new identity satisfied for every bounded solution of the nonlocal semilinear Dirichlet problem $(-\Delta)^s u = f(x, u)$ in Ω , $u \equiv 0$ in $\mathbb{R}^n \setminus \Omega$. Here, $s \in (0, 1)$, and $(-\Delta)^s$ stands for the fractional Laplacian in \mathbb{R}^n —the infinitesimal generator of a symmetric and stable Lévy process. Surprisingly, from a nonlocal problem we obtain an identity with a boundary term (an integral over $\partial\Omega$) which is completely local.

To establish this identity we need to prove, among other things, that if u is a bounded solution then $u/d^s|_{\Omega}$ is C^α up to the boundary $\partial\Omega$, where $d(x) = \text{dist}(x, \partial\Omega)$. In our Pohozaev identity, the function $u/d^s|_{\partial\Omega}$ plays the role that $\partial u/\partial\nu$ plays in the classical one.

LUGAR / LEKUA:

Sala de seminarios de la sección de matemáticas
Matematika ataleko mintegi gela

DÍA Y HORA / EGUNA ETA ORDUA:

29/05/2014, 11:00